ELSEVIER

Carbohydrate Research 288 (1996) C7–C8

CARBOHYDRATE RESEARCH

Contents

Regular Papers

Cross-polarisation kinetics and proton NMR relaxation in polymers of *Citrus* cell walls
M.C. Jarvis, K.M. Fenwick (Glasgow, UK) and D.C. Apperley (Durham, UK) 1

CP-MAS NMR of highly mobile hydrated biopolymers: polysaccharides of *Allium* cell walls
M.-A. Ha, B.W. Evans, M.C. Jarvis (Glasgow, UK), D.C. Apperley and A.M. Kenwright (Durham, UK) 15

Synthesis and structural studies of branched 2-linked trisaccharides related to blood group determinants
J.Ø. Duus (Copenhagen, Denmark), N. Nifant'ev, A.S. Shashkov, E.A. Khatuntseva (Moscow, Russia) and K. Bock (Copenhagen, Denmark) 25

Mutarotation of D-fructose in aqueous–ethanolic solutions and its influence on crystallisation
A.E. Flood, M.R. Johns and E.T. White (Brisbane, Australia) 45

Predicting helical structures of the exopolysaccharide produced by *Lactobacillus sake* 0-1
G.W. Robijn (Utrecht, The Netherlands), A. Imberty (Nantes, France), D.J.C. van den Berg, A.M. Ledeboer (Vlaardingen, The Netherlands), J.P. Kamerling, J.F.G. Vliegenthart (Utrecht, The Netherlands) and S. Pérez (Nantes, France) 57

Visible and ^{13}C nuclear magnetic resonance spectra of azo dyes and their complexes with cyclomalto-oligosaccharides
M. Suzuki, H. Takai, K. Tanaka, K. Narita, H. Fujiwara and H. Ohmori (Osaka, Japan) 75

Synthesis of four glycosides of a disaccharide fragment representing the terminus of the O-polysaccharide of *Vibrio cholerae* O:1, serotype Inaba, bearing aglycons suitable for linking to proteins
Y. Ogawa, P.-s. Lei and P. Kováč (Bethesda, MD, USA) 85

Synthesis of a tetrahydropyrano[2,3-*d*]oxazole analogue of trehazolin
M. Shiozaki, T. Mochizuki, H. Hanzawa and H. Haruyama (Tokyo, Japan) 99

The use of 2-deoxy-2-trichloroacetamido-D-glucopyranose derivatives in syntheses of hyaluronic acid-related tetra-, hexa-, and octa-saccharides having a methyl β-D-glucopyranosiduronic acid at the reducing end
G. Blatter and J.-C. Jacquinet (Orléans, France) 109

A novel enzymic determination of maltobionate
Y. Shirokane, A. Arai, R. Uchida and M. Suzuki (Noda City, Japan) 127

Characterisation of acidic heteroglycans from *Tremella fuciformis* Berk with cytokine stimulating activity
Q. Gao, R. Seljelid (Tromsø, Norway), H. Chen and R. Jiang (Changchun, People's Republic of China) 135

Arabinose and ferulic acid rich pectic polysaccharides extracted from sugar beet pulp
A. Oosterveld, G. Beldman, H.A. Schols and A.G.J. Voragen (Wageningen, The Netherlands) . . . 143

Composition and structural analysis of alpha-dextrins from potato amylopectin
Q. Zhu and E. Bertoft (Turku, Finland) 155

The influence of the conformational state of κ- and ι-carrageenan on the rate of acid hydrolysis
T. Hjerde, O. Smidsrød and B.E. Christensen (Trondheim, Norway). 175

Structure of stewartan, the capsular exopolysaccharide from the corn pathogen *Erwinia stewartii*
M. Nimtz (Braunschweig, Germany), A. Mort (Stillwater, OK, USA), V. Wray, T. Domke (Braunschweig, Germany), Y. Zhang (Heidelberg, Germany), D.L. Coplin (Columbus, OH, USA) and K. Geider (Heidelberg, Germany). 189

Structural characterization of the exopolysaccharide produced by *Lactobacillus acidophilus* LMG9433
G.W. Robijn, R.G. Gallego (Utrecht, The Netherlands), D.J.C. van den Berg, H. Haas (Vlaardingen, The Netherlands), J.P. Kamerling and J.F.G. Vliegenthart (Utrecht, Netherlands) 203

Potato xyloglucan is built from XXGG-type subunits
J.-P. Vincken, A.J.M. Wijsman, G. Beldman (Wageningen, The Netherlands), W.M.A. Niessen (Leiden, The Netherlands) and A.G.J. Voragen (Wageningen, The Netherlands) 219

Notes

Reaction of enzymes with starch granules: enhanced reaction of glucoamylase with gelatinized starch granules
A. Kimura and J.F. Robyt (Ames, IA, USA) 233

Stereospecific synthesis of (−)-*allo*-muscarine from D-glucose
V. Popsavin, O. Berić, M. Popsavin, J. Csanádi and D. Miljković (Novi Sad, Yugoslavia) 241

Author index C1

Subject index C3

Contents C7